

UNIVERSITY OF BIRMINGHAM

School of Computer Science

Mathematical and Logical Foundations of Computer Science

First Class Test 2021/22

This test is designed to be solved in about one hour and is worth 7% of your total grade.

The module was taught differently last year.
Only attempt the questions circled in red,
not the striked out ones.

Mathematical and Logical Foundations of Computer Science

Question 1 [Numbers and Set Theory]

- (a) (i) Suppose A, B, C are subsets of some set X . Draw a Venn diagram for the expression $(A \setminus B) \cup ((B \cap C) \setminus A)$. [2 marks]
- (ii) Find an expression for $(A \setminus B) \cup ((B \cap C) \setminus A)$ that uses only union, intersection, and complement. [2 marks]
- (iii) For the sets $D = \{x \in \mathbb{Z} \mid x^2 \leq 40\}$ and $E = \{x \in \mathbb{Z} \mid \text{there exists } y \in \mathbb{Z} \text{ such that } 3y = x\}$ write down D and $D \cap E$ explicitly. [4 marks]
- (b) (i) Does $\mathbb{Z}_6$ satisfy the law of the multiplicative inverse? In other words, for each $x \in \mathbb{Z}_6$ does there exist $y \in \mathbb{Z}_6$ such that $xy \equiv 1 \pmod{6}$? Justify your answer.
assume that x is not zero for the question above [2 marks]
- (ii) Consider the following piece of pseudocode, where n is a natural number:
- ```
x ← 0
s ← 0
while (x < n) {
 x ← x + 1
 s ← s + 2*x - 1
}
return s
```
- Prove that  $s = x^2$  is an invariant of the loop. [4 marks]
- (c) Java offers functionality for creating arrays of floating point numbers, which have the type `float[]`. The length of such an array is specified as an `int` variable. Consider the set of all possible arrays of type `float[]`. Is the cardinality of this set finite, countable, or uncountable? Discuss the relationship of this set to the sets of lists and streams of numbers that we defined. [6 marks]

## Question 2 [Propositional Logic]

- (a) Let  $F$  be the following proposition:  $(\neg A \vee \neg B) \rightarrow (C \rightarrow A \wedge B) \rightarrow \neg C$ . Provide an intuitionistic Natural Deduction proof of  $F$ . [8 marks]
- (b) Let  $G$  be the following proposition:  $\neg\neg\neg A \rightarrow \neg A$ .
- (i) Provide an intuitionistic Natural Deduction proof of  $G$  [4 marks]
- (ii) Provide a intuitionistic Sequent Calculus proof of  $G$ . [4 marks]
- (c) Let  $H$  be  $(\neg P \rightarrow Q \wedge R) \rightarrow P \vee R$ . Provide a proof of  $H$  using the 2nd classical version of the Sequent Calculus (i.e. the version without additional LEM or DNE rules but with classical sequents instead). [4 marks]